

Ergasiophytes of the Ukrainian flora

Vira V. Protopopova & Myroslav V. Shevera*

M. G. Kholodny Institute of Botany, National Academy of Sciences of Ukraine, Tereschenkivska 2, 01601 Kiev, Ukraine
* corresponding author (e-mail: shevera@mail.ru)

Abstract: The results of an investigation of the ergasiophytes (species deliberately introduced to a region for cultivation) of the Ukraine (458 species) are presented. The systematic, life forms and morphological types of plants with respect to soil moisture spectras, origin and degree of naturalisation of species of this group are analyzed. A annotated list of ergasiophytes is provided.

Key words: alien species, ergasiophytes, invasive plants, Ukraine

1. Introduction

By the end of the 20th century, invasions of alien organisms, including plants, were widely realized as one of the major global threats to biodiversity (Pyšek *et al.* 1995; Baldacchino & Pizzuto 1996; Mooney & Cleland 2001; Reichard & White 2001; Kowarik 2002; Protopopova *et al.* 2002, 2003; Chornesky & Randall 2003; Davis 2003; Sax & Gaines 2003, etc.).

Ergasiophytes (species deliberately introduced to a region for cultivation) play a significant role synanthropisation, the process by which the native flora of a region becomes replaced by alien species. These species generally are delivered into a new region from places where they were cultivated. Therefore they are partially adapted to condition of anthropogenic habitats. So these species can adapt more easy and quickly to new conditions. The detection of the new species of alien invasive plants before they become invasive, with the aim of preventing their uncontrolled spread is one of necessary conditions of the *Convention on Biological Diversity* (1994), the *Global strategy on invasive alien species* (2001), the *European strategy on invasive alien species* (Genovesi & Shine 2004) and other international programmes and documents. It is the responsibility of every country which signed them, including Ukraine.

In compiling the list of the Ukrainian alien fraction flora has drawn attention to the increasing role of species which have been intentional introduction into the culture and later escaped. The number of ergasiophytes

increases and many of the species are characterized by a high degree of naturalization. Therefore the necessity of special complex study of ergasiophytes arises.

The main aim of the study was to clarify of modern species composition of the Ukrainian flora ergasiophytes and their some characteristics.

The objectives of the study included: 1) inventory of ergasiophytes species, and preparation of the annotated list; 2) generalization of their systematic, biological and ecological characteristics, origin, as well as determination of the degree of naturalization in the region.

2. Material and methods

The present research focuses on the ergasiophytes of the Ukrainian alien fraction flora. The investigations are based on original data obtained by routine surveys in 2010-2014, the analysis of data in the literature and examinations of the collections of the Herbaria of M. G. Kholodny Institute of Botany, National Academy of Sciences of Ukraine (KW), Donetsk Botanical Garden, NAS of Ukraine (DNZ), Uzhgorod National University (UU), Y. Fedykovich National University (CHER), I. Franko Lviv National University (LW), T. Shevchenko Kyiv National University (KNU), I. I. Mechnikov Odessa National University (MSDU), V. N. Karazin Kharkiv National University (CWU), V. I. Vernadskyi Tavrian National University (SIMF), Crimean Branch of Agrotechnology Institute of National University of Biological Resources and Natural Uses (CSAU), Nikita

Botanical Garden-National Science Centre, Ukrainian Agricultural Academy of Sciences (YALT).

Comparative morphological and geographical methods were used in the present investigations. Taxonomic structure (according to Tolmachev 1974), life forms (according to Serebryakov 1962), and morphological types of plants with respect to soil moisture (according to Didukh 2000) of the ergasiophytes were analysed. Characteristics of area species prepared according to Flora of European part of USSR and Flora of Eastern Europe (Fedorov 1974-1987; Tzvelev 1989-1994, 1996-2004). The classification of alien plants proposed by Kornaś and modified by Protopopova (1991) and Moysiенко (1999) was used, e.g. ‘agrio-epoecophyte’ – species which fully naturalized in anthropogenic, semi-natural and natural plant communities. Invasive and transformer groups determined according Richardson *et al.* (2000). The species names are given according to Mosyakin & Fedorochuk (1999).

3. Results

Studies of alien plants over 25 year period testified that the process of synanthropisation of the floras of natural zones (Forest, Forest-Steppe, and Steppe and their divisions) and in the Ukrainian flora in general has been constantly progressing. We can observe clear tendencies for the number of alien species to increase, the growth of both ‘naturalised’ or ‘stable’ (agriophytes, agrio-epoecophytes, epoecophytes, colonophytes) and ‘unstable’ (ephemeralophytes, ergasiophytes) components of the alien fraction of the flora, growing rates of immigration and dispersal, an increasing number and scope of invasions of certain invasive species (Protopopova *et al.* 2002, 2003).

The ergasiophyte group of the Ukraine, according to our data, is represented by at least 458 species of

vascular plants from 297 genera and 92 families (see Appendix 1), or 46.3% of the alien fraction flora and 7.5% of total number of the Ukrainian flora. Tendency of increasing number of the Ukrainian alien fraction flora (black line) and ergasiophytes group species (gray line) during 150 years is presented on Fig. 1.

The taxonomic spectrum of leading places of ergasiophytes of the Ukraine includes 12 families: Asteraceae (59 sp.), Rosaceae (36), Fabaceae (34), Brassicaceae and Poaceae (24 each), Apiaceae (16) Lamiaceae (14), Solanaceae (13), Caryophyllaceae (12), Polygonaceae (10), Chenopodiaceae and Malvaceae (9 each). These leading families comprise 260 species or 56.8% of the total. Three of leading families comprise 130 species or 28.6%. Other families contain from eight to one species, including 28 families which are presented only one or two species. The taxonomic spectrum of the leading families of ergasiophytes differs from the general spectra of the Ukrainian alien fraction flora as does the relative position of the families (excluding Asteraceae) and the presence of new ones (Rosaceae, Solanaceae, Polygonaceae).

The taxonomic spectrum of the leading genera of studied group is not clearly expressed. The largest number of species (7) are from the genus *Helianthus* L. The genera *Opuntia* Mill. and *Sympyotrichum* Nees include six species each; *Acer* L., *Allium* L., *Amaranthus* L. and *Brassica* L. – five species each; *Datura* L., *Juglans* L., *Lonicera* L., *Populus* L., *Prunus* L. *Pyrethrum* Zinn, *Rosa* L., *Spiraea* L., *Trifolium* L. and *Vitis* L. – four species each; 26 genera include three species each, 40 genera – two species each, and 214 – only one species each.

In the spectra of life forms of species of ergasiophytes of Ukraine (Fig. 2), as in the alien fraction flora, annuals prevail (142 sp. or 31%), but their participation percentage is lower. The second position is occupied by




Fig. 1. Dynamics of the number of the Ukrainian alien fraction flora species (black line) and the Ukrainian ergasiophytes (gray line)




Fig. 2. Spectrum of life form of species of ergasiophytes of the Ukrainian flora

herbaceous perennials (134 or 29.3%). A peculiarity of the spectra is the high participation of the shrubs and trees, among them there are 58 shrubs, 51 trees, 23 shrub or tree, 2 semishrubs (in total 133 species or 29.3%). Other groups do not include numerous species and are represented by from three to one species.

Among the ergasiophyte species of the Ukrainian flora with respect to soil moisture (Fig. 3) xero-mesophytes prevail (196 species or 42.8%). The second position occupied mesophytes (176 or 38.4%), the next

position – meso-xerophytes (41 or 8.9%) and xerophytes (31 or 6.7%). The other groups, hydrophytes, hygrophytes, hygro-mesophytes and hydro-mesophytes, are unnumerous and presented from six to one species.

The species of ergasiophytes of the Ukrainian flora vary in origin. With regard to their primary area of origin (Fedorov 1974-1987; Tzvelev 1989-1994, 1996-2004) ergasiophytes are divided into a number of types. The basis for this group is presented by origin from North America (105 species), the Mediterranean (85),




Fig. 3. Spectrum of the morphological types of plants with respect to soil moisture of species of ergasiophytes of the Ukrainian flora

East Asia and Asia (26 each). The next large groups are South American (16), South European (15), Caucasic (13), West Mediterranean (11) and West European (10) in origin. The group of hybrid origin is represented by 16 species. Other groups do not include numerous species and are represented by from nine to one species, including 27 types of area and subdivision which consist one species only.

With respect to the time of immigration, the species of ergasiophytes of Ukrainian flora are divided into archaeophytes (9 species) and kenophytes (449). Some species have a different status in different natural zones of Ukraine, e.g. the species *Acer tataricum* L. which is native in Forest-Steppe zones or *Rubus idaeus* L., native in Forest zones, are kenophytes in Crimea.

With regard to their degree of naturalization, the species of ergasiophytes of the Ukrainian flora may be divided into the 7 groups. e.g. agriophytes (27), agrio-epoecophytes (21), agrio-colonophytes (1), epoecophytes (22), colonophytes (66), ephemeralophytes (43) and ergasiophygophytes (278). The degree of naturalization of numerous species differs in the various botanical and geographical regions of the Ukraine, e.g., *Antirrhinum majus* L. is an ergasiophygophyte in major of regions of Ukraine and an agriophyte in Crimea.

The extent of distribution of species of ergasiophytes varies. The distribution of a large part of the ephemeralophytes and ergasiophygophytes has a pattern of a single or a few localities; some of them however were cultivated over a long period of time and are very common and sporadically distributed throughout the territory of the Ukraine. The majority of epoecophytes and colonophytes are distributed in more than three botanical and geographical regions of the Ukraine. Agriophytes are a component of plant communities in one or two regions. A tendency to active spread is most evident for agrio-epoecophytes and epoecophytes. The distribution of agriophytes is limited by the presence of corresponding plant communities susceptible to invasion. Thus, from the 458 species of ergasiophytes 137 or 29.9% were considered to be completely naturalized. Among them we have identified 31 species or 6.7% which may be defined as invasive, including 17 species or 3.7% transformers.

The cultivation of plants in Ukraine has an ancient history. The primary centres of cultivation were the monasteries which cultivated mainly medicinal plants, later apothecary gardens and private parks where numerous ornamental plants were planted. One of the first botanical gardens were created in Lubny (1721), Kharkiv (1804), Kremenets (1806), Nikita (1812), Odesa and Nizhyn (1820). Some alien species which were cultivated there now are widely distributed in Ukraine, e.g. *Cyclachaena xanthiiifolia* (Nutt.) Fresen. (from Kiev University Botanical Garden), *Ailanthus altissima* (Mill.) Swingle and *Bupleurum fruticosum* L. (from Nikita Botanical Garden, Crimea), *Acer negundo*

L. and *Ailanthus altissima* (from Osnovyanskyi, modern Krasnokutskyi, Park, near Kharkiv), *Robinia pseudoacacia* L. (from Odesa and Kharkiv Botanical Gardens), *Abutilon theophrasti* Medik. (from Lubny Botanical Garden, Poltava Reg.), etc. In the 20th century an essential role in the cultivation of new plants for agricultural and technical purposes was taken up by Experimental Stations (Kharkiv, Poltava, Lubny, Maslivka, etc.). In recent years the uncontrolled import of wide range of ornamental and vegetable plants seed has been observed.

We have attempted to reconstruct the main stages of naturalization of some ergasiophytes (Table 1).

For the most part of this group, data about their first cultivation are absent. The literature (Palimpsestov 1855; Protopopova 1973; Vynogradova *et al.* 2010; Mayorov *et al.* 2012, etc.), Herbaria materials, and original data concerning the naturalization of transformer species shows that the period from the first note as escaped plants to their expansion is from over 20 years to 100 or more, e.g. *Impatiens parviflora* DC. was known in cultivation in Ukraine from 1871 in Lviv gardens and parks, and from 1895 – in Dublyany park (Lviv Region). About 40 years later (1908) the first escaped plants of this species were noted in the Lviv area and in the Carpathians. The active distribution of this species was beginning 50 years ago and continues at present. In Europe as a whole, this period has lasted 50 years longer. Now in Polissya (the Forest zone) the species has been noted in plant communities of the *Querco-Fagetea* Br.-Bl. et Vlieg. 1937, *Robinieta Jurko ex Hadač et Sofron* 1980 and *Galio-Urticetea* Pass. 1967 em. Kopecky 1969 classes (Lukash 2008), and in Forest-Steppe zones of the *Alnetea glutinosae* Br.-Bl. et R. Tx. 1943 class also (Kaniv State Reserve) and in Bukovyna Cis-Carpathian in *Epilobietea angustifolii* Tx. et Prsg. ex von Rochow 1951, *Artemisieta vulgaris* Lohm., Prsg et R. Tx. in R. Tx. 1950, *Polygono-Poetea annuae* Rivas-Mart. 1975, and *Galio-Urticetea*, and in *Querco-Fagetea*, *Robinieta* classes (Protopopova *et al.* 2010). A similar situation has been observed for *Impatiens glandulifera* Royle. Presently this species in Polissya is a component of plant communities of the *Phragmito-Magnocaricetea* Klika in Klika et Novak 1941, *Molinio-Arrhenatheretea* R. Tx. 1937, *Alnetea glutinosae*, *Bidentea tripartiti* R. Tx., Lohm. et Prsg 1950 classes. In *Echinocystis lobata* (Michx.) Torr. & A. Gray the period of adaptation was about 40 years in Ukraine and 30-80 – in other parts of Europe. The period from the beginning of cultivation to an expansion in *Heracleum sosnowskyi* Manden was about 20 years, in *Solidago canadensis* L. and *Senecio cineraria* DC. – about 100 (Protopopova *et al.* 2000, 2012; Protopopova & Shevera 2013a, 2013b; Vinogradova *et al.* 2010). These examples show the different rates at which some species adapt to new conditions.

Now ergasiophytes are registered in Ukraine as components of almost all types of semi-natural and natural

Table 1. The main stages of naturalization of ergasiophytes in Ukraine

Species	First date of cultivation in Europe, country	First date of cultivation in Ukraine	Date and region of first registration as escaped plants	The start of active distribution and region
<i>Acer negundo</i>	1688, England	1809, Osnovyanskyi (modern Krasnokutskyi) dendrological park (Kharkiv Reg.); 1814 – Nikita Bot. Gard. (Crimea); 1816 – Kremenets Bot. Gard (Ternopil Reg.); 1825-1830 – Trykratskyi park of Skarzhynskyi (modern Odesa Reg.); 1865 – Velyko-Anadolskyi forestry (modern Donetsk Reg.)	1850s-1860s, ?	The second half of 20 th c. Forest-Steppe zone
<i>Ailanthus altissima</i>	1740, ?	1809, Osnovyanskyi; 1814 – Nikita Bot. Gard.; 1820 – Simferopol, Sebastopol, Theodosia	1835, Crimea	The end of 20 th c., Crimea
<i>Ambrosia artemisiifolia</i>	1863, Germany	1914, vill. Kudashivka (modern Dnipropetrovsk Reg.)	1925, Kiev	1950s-1960s, Steppe zone
<i>Amorpha fruticosa</i>	1724, England	1930s or middle of 19 th c., Odesa parks, and Trykratsky park of Skarzhynsky (Odesa Reg.)	Probably the middle of 20 th c., ?	1990s, Middle Dnipro Region
<i>Bupleurum fruticosum</i>	-	1814, Nikita Bot. Gard.	1885, Alupka (Crimea)	1914, South-Western Crimea
<i>Echinocystis lobata</i>	The second half of 19 th c., ?	Probably first half of 20 th c.	1933, vill. Didivtsi, Transcarpathia	1980s, western regions of Ukraine
<i>Elaeagnus angustifolia</i>	1792, Romania	1930s, Odesa gardens	1925, vill. Foros and Sudak (Crimea)	1990s, steppe regions of Ukraine
<i>Heracleum mantegazzianum</i>	1817, England	1927, Osmoloda forestry (modern Ivano-Frankivsk Reg.)	1962, vill. Osmoloda	1990s, Transcarpathia
<i>Heracleum sosnowskyi</i>	1947, Russia	1960s-1970s, western and northern forest regions of Ukraine	-	90 years of 20 th c., Transcarpathia, Polissya (Forest zone)
<i>Impatiens parviflora</i>	1831, Switzerland	1871, Lviv parks	1908, Lviv	The middle of 20 th c. Forest-Steppe zone and Polissya (Forest zone)
<i>Impatiens glandulifera</i>	1838, England	1930s, Transcarpathia and some western regions of Ukraine	1938, vill. Osiy and Hankovytsha, Transcarpathia; 1939, vill. Mykhaylivka, Khmelnytskyi Reg.	1990s, Transcarpathia
<i>Helianthus tuberosus</i>	1774, ?	?	?	1990s, Transcarpathia
<i>Reynoutria japonica</i>	1825, England	?	1929, Rachiv, Transcarpathia	1990s, Transcarpathia
<i>Rhamnus alaternus</i>		1812, Nikita Bot. Gard.	1860	Probably 1950s, Crimea (South bank)
<i>Rudbeckia laciniata</i>	1663, ?	the middle (?) and the end of 19 th c. – the beginning of 20 th c., Transcarpathia, Bukovyna	1903, vicinity of vill. Serednye, Transcarpathia, 1911 vill. Jordaneshti, Bykovyna	The end of the 20 th c. Bukovyna, Transcarpathia
<i>Senecio cineraria</i>	?	First half of 19 th c., probably in Nikita Bot. Gard.	1842	1950s Crimea (South bank)
<i>Solidago canadensis</i>	1645, England	The beginning of 19 th c.	1886	1990s right bank of Forest-Steppe zone

plant communities and ecosystems (forest, steppe, aquatic vegetation, etc.), e.g. in forests we registered invasions of *Acer negundo*, *Impatiens parviflora*; along river shores, *Heracleum mantegazzianum*, *Impatiens glandulifera*, *Echinocystis lobata*, *Reynoutria japonica*, and several species of *Helianthus* L. form large, often monodominant, populations. The dispersal of invasive species represents a serious danger to native rare species and natural protected areas, e.g. the most dangerous alien plant in the Danube Biosphere Reserve in sandy habitats – *Elaeagnus angustifolium*, in Kaniv State Reserve in flood plain forest – *Acer negundo* L. and *Amorpha fruticosa* L. (Protopopova *et al.* 2006a).

4. Discussion

The spectrum of ergasiophytes according to origin of species are differentiated by large variety, however the success of species naturalization persists in those groups that in the general range of the Ukrainian alien fraction flora. The greatest success has a species of the North American and East Asian origin. As in the general spectrum of the Ukrainian alien fraction flora among ergasiophytes the species of Mediterranean origin are dominated, however most of them have a low degree of naturalization. Feature of the life form spectrum is higher percentage of perennials, shrubs and trees among the ergasiophytes; and in the morphological types of plants with respect to soil moisture spectrum the mesophytes group (in an aggregate) is dominated compared with the general spectrum of the Ukrainian alien fraction flora.

Growing of the species in culture at the first stages under the conditions of new region facilitate the naturalization of plants. In the literature (Vynogradova *et al.* 2010) the opinion was expressed that the seed and planting material are often imported not from primary habitats, and of the places where these species are cultivated. And that means they are already partly preadapted to the culture conditions. Nowadays this is also evidenced by reducing the time of naturalization of certain species. Increasing the number and diversity of anthropogenic and disturbed natural ecotopes that contribute to the emergence of new niches facilitates the introduction of alien species into synanthropic and disturbed natural plant communities. For example, in Ukraine due to changes of climatic conditions the mesophytisation

of the Steppe zone ecotopes is observed, the changing of their water and salt regimes is noted (Vyhnevsky 2000). It has caused suppression of several species of local flora and facilitated naturalization of some ergasiophytes adapted to the conditions which have arisen. For example, *Elaeagnus angustifolia* actively consumes water resources and overcompetes many local species; it is a salt-tolerant species. It forms stands that change the light mode of the herbaceous vegetation layer in Steppe zone communities, thus affecting the species composition and the structure of plant communities. This species forms the association *Elaeagnetum angustifoliae* Chinkina, and occupies specific econiches in other communities, e.g., *Hippophaeta rhamnoidis*, thus promoting degradation of aboriginal communities. The species *E. angustifolia* shows high invasive ability to penetrate many different types of habitats (ecotopes) and plant communities, which occurs in different floristic complexes (natural: psammophyton, pratophyton, halophyton, steppophyton, drymophyton, petrophyton, litoralophyton, and anthropogenic: aggeratophyton, and runkatiodyrmophyton). The species occurs mostly on sandy and riverside semi-natural ecotopes with unstable and sparse plant cover. The most diversity of plant communities with participation of *E. angustifolia* is observed in sandy (8 associations) and riverside (7 associations) biotopes with sparse vegetation. The main limiting factors for the species distribution in the studied region are excessively humid ecotopes (Protopopova *et al.* 2006, 2009).

5. Conclusion

The above data shows that, the ergasiophyte group, represented by 458 species in Ukraine, plays a significant role in the enrichment of spontaneous flora (Protopopova & Shevera 2013a, 2013b). The results of our studies show that numerous species of this group reached a high degree of naturalization and are widely distributed.

Acknowledgement. We would like to thank Ms. Natalia V. Morgart (M. G. Kholodny Institute of Botany, NAS of Ukraine) for technical assistance. The authors are grateful to an anonymous reviewer for his detailed review of manuscript, valuable comments and suggestions.

References

- BAGRIKOVA N. A. 2013. Structural analysis of the alien fraction of the flora of the Crimean peninsula (Ukraine). Ukr. Bot. Journ. 70 (4): 489-507.
- BALDACCHINO A. E. & PIZZUTO A. (eds.). 1996. Introduction of alien species of flora and fauna. Proceedings of Seminar held at Qawra, Malta, 5th March 1996. p. 77. Malta: EPD-Floriana.
- CHORNESKY E. A. & RANDALL J.M. 2003. The threat of invasive alien species to biological diversity: Setting a future course. Ann. Missouri Bot. Gard. 90(1): 67-76.
- Convention on Biological Diversity. 1994. Text and Annexes. 34 pp. Chatelaine, Switzerland: UNEP/CBD.
- DAVIS M. 2003. Biotic Globalization: does competition from introduced species threaten biodiversity? BioScience 53: 481-489.
- DIDUKH YA. P. (ed.). 2000. Ecoflora of Ukraine. Vol. 1. 284 pp. Fitosociocenter Press, Kiev.
- FEDOROV AN. A. (ed.). 1974-1987. Flora of European part of the USSR. Vol. 1-6. Nauka Press, Leningrad.
- GENOVESI P. & SHINE C. 2004. European strategy on invasive alien species. 68 pp. Council of Europe, Strasbourg.
- Global Strategy on Invasive Alien Species / Convention of Biological Diversity, 2001. SBSTTA Sixth Meeting. Montreal, ix+52 pp.
- KORNAŚ J. 1968. Geograficzno-historyczna klasyfikacja roślin synantropijnych. Mater. Zakł. Fitosoc. Stos. U.W. 25: 33-41.
- KOWARIK I. 2002. Biologische Invasionen in Deutschland: zur Rolle nichtheimischer Pflanzen. In: I. KOWARIK & U. STARFINGER (eds.). Biologische Invasionen. Herausforderung zum Handeln. Neobiota 1: 5-24.
- LUKASH O. V. 2008. Flora of vascular plants of Eastern Polissya: history of investigation, conspect. 436 pp. Phytosociocenter Press, Kiev.
- MAYOROV S. R., BOCHKIN V. D., NASYMOVICH YU. A. & SCHERBAKOV A. V. 2012. Alien flora of Moscow and Moscow Region. 412 pp. Tovarischestvo nauchnykh izdaniy KMK Press, Moscow.
- MOYSIYENKO I. I. 1999. Urban flora of Kherson. Comprehensive summary of PhD Thesis. Yalta. 19 pp.
- MOONEY H. A. & CLELAND E. E. 2001. The evolutionary impact of invasive species. Proceeding NAS USA. 98: 5446-5451.
- MOSYAKIN S.L. & FEDORONCHUK N. M. 1999. Vascular plants of Ukraine. A nomenclatural checklist. xxiii+345 pp. Institute of Botany, NAS of Ukraine, Kiev.
- Ostapko V. M., Boyko A. V. & Mosyakin S. L. 2010. Vascular plants of South-Eastern of Ukraine. 247 pp. Noulidzh Press, Donetsk.
- PALIMPSESTOV N. 1855. Vocabulary of agricultural plants. 790 pp. Odessa.
- PROTOPOPOVA V. V. 1973. The adventive species of Forest-Steppe and Steppe of Ukraine. 192 pp. Naukova Dumka Press, Kiev.
- PROTOPOPOVA V. V. 1991. The Synanthropic Flora of Ukraine and its Development. 204 pp. Naukova Dumka Press, Kiev.
- PROTOPOPOVA V. V., MOSYAKIN S. L. & SHEVERA M. V. 2002. Plant invasions in Ukraine as a threat to biodiversity: The present situation and tasks for the future. 32 pp. M. G. Kholodny Institute of Botany, NAS of Ukraine, Kiev.
- PROTOPOPOVA V. V., MOSYAKIN S. L. & SHEVERA M. V. 2003. Impact of alien plant species on the phytobiota of Ukraine. In: O. V. DUDKIN (ed.). Assessment and Mitigation of Threats to Biodiversity of Ukraine, pp. 129-155. Chimjest Publ., Kiev.
- PROTOPOPOVA V. V. & SHEVERA M. V. 2013a. Ergasiophytes in Ukrainian flora: present state and degree of risk. The role of botanical gardens and dendrological parks in conservation and enrichment of biological diversity of urban territories. In: V. RADCHENKO (ed.). Proceeding of the International sciences conference, Kiev, 2013, May 28-31, pp. 138-139. Vitpol Press. Kiev.
- PROTOPOPOVA V. V. & SHEVERA M. V. 2013b. Ergasiophytes as a potential reserve of Ukrainian alien fraction flora. Non-traditional, new, and forgotten species of plants: sciences and practical aspects of cultivation. Proceeding of the International sciences conference, Kiev, 2013, Sept. 10-12), pp. 99-101. Knyganosha Press, Kiev.
- PROTOPOPOVA V. V., SHEVERA M. V., BAGRIKOVA N. A. & RYFF L. E. 2012. Transformer species in the flora of the South Coast of Crimea. Ukr. Bot. Journ. 69(1): 54-68.
- PROTOPOPOVA V. V. SHEVERA M. V., CHORNEY I. I., TOKARYUK A. I., BUDZHAK V. V. & KORZHAN K. V. 2010. The transformer species in the flora of the Bukovyna Cis-Carpathian area. Ukr. Bot. Journ. 67(6): 864-862.
- PROTOPOPOVA V. V., SHEVERA M. V. & MELNYK R. P. 2006. The history of introduction and present distribution of *Elaeagnus angustifolia* L. in the Black Sea Region of Ukraine. Chornomorski Bot. Journ. 2(2): 5-13.
- PROTOPOPOVA V. V., SHEVERA M. V. & MOSYAKIN S. L. 2006. Deliberate and unintentional introduction of invasive weeds: a case study of the alien flora of Ukraine. Euphytica 148: 17-33.
- PYŠEK P., DANIELKA J., SÁDLO J., CHRTEK J. JR., CHYTRÝ M., JAROŠÍK V., KAPLAN Z., KRAHULEC F., MORAVCOVÁ L., PERGL J., ŠTAJEROVÁ K. & TICHÝ L. 2012. Catalogue of alien plants of the Czech Republic (2nd edition): checklist update, taxonomic diversity and invasion patterns. Preslia 84: 155-255.
- PYŠEK P., PRACH K., REJMÁNEK M. & WADE M. (eds.). 1995. Plant Invasions. General Aspects and Special Problems. 257 pp. SPB Academic Publishing, Amsterdam.
- REICHARD S. H. & WHITE P. S. 2001. Horticulture as a pathway of invasive plant introductions in the United States. BioScience 51: 103-113.
- REICHARD, S. H. & WHITE P. S. 2003. Invasion biology: An emerging field of study. Ann Missouri Bot Gard 90: 64-66.
- RICHARDSON D. M., PYŠEK P., REJMÁNEK M., BARBOUR M. G., PANETTA D. D. & WEST C. J. 2000. Diversity Distrib. 6: 93-107.

- SAX D. F. & GAINES S. D. 2003. Species diversity: From global decreases to local increases. *Trends Ecol. Evol.* 18: 541-545.
- SEREBRYAKOV I. G. 1962. The ecological morphology of plants. Life forms of angiosperms and gymnosperms. 379 pp. Vischaya shkola Press, Moskow.
- TOLMACHEV A. I. 1974. Introduction in the geography of plants. 244 pp. Leningrad State University Press, Leningrad.
- TZVELEV N. N. (ed.). 1989-1994. Flora of European part of the USSR. Vol. 7-8. Nauka Press, Leningrad.
- TZVELEV N. N. (ed.). 1996-2004. Flora of Eastern Europe. Vol. 9-11. Mir i Semya-1995 Press and Tovarischestvo nauchnykh izdaniy KMK Press. St.-Peterburg, Moscow.
- VYSHNEVSKY V. I. 2000. The Rivers and reservoirs of Ukraine. State and use. 376 pp. Vipol Press, Kiev.

Appendix 1. Annotation list of ergasiophytes of the Ukrainian flora

The list including the next positions: name of species, family, life form, hydromorphe, chronological elements (data of first publication or fixation), origin, degree of naturalization, way of distribution; invasive or transformer (region).

- Abutilon theophrasti* Medik., Malvaceae, Ann., Xero-meso, Ken. (1855), E. As., Epoec., Erg.
- Acer negundo* L., Aceraceae, Tree, Meso, Ken., N. Am., Agr.-epoec., Erg.-kseno; invasive; transformer (forest and forest-steppe regions of Ukraine).
- Acer platanoides* L., Aceraceae, Tree, Meso, Ken. (Crimea)/natural species in Ukraine, Eur., Col., Erg.
- Acer pseudoplatanus* L., Aceraceae, Tree, Meso, Ken. (Crimea)/natural species in Ukraine, Eur.-Cauc. (Tzvelev 1996), Col., Erg.
- Acer saccharinum* L., Aceraceae, Tree, Meso, Ken., N. Am., Ergasphig., Erg.
- Acer tataricum* L., Aceraceae, Tree, Xero-meso, Ken. (Crimea)/natural species in Ukraine, Eur., Col., Erg.
- Acorus calamus* L., Araceae, Perenn., Hygro, Arch., As., Agr., Erg.; invasive.
- Aesculus hippocastanum* L., Hippocastanaceae, Trees, Meso, Ken., Med., Ergasphig., Erg.
- Agrostemma githago* L., Caryophyllaceae, Ann., Xero-meso, Arch., Anthroc., Ergasphig., Kseno-erg.
- Ailanthus altissima* (Mill.) Swingle, Simaroubaceae, Shrub, Xero-meso, Ken., E. As., Agr.-epoec., Erg.; invasive; transformer (Crimea).
- Albizia julibrissin* Durazz., Mimosaceae, Tree, Xero-meso, Ken. (Crimea), S. E. Transcauc., Col., Erg.
- Alcea rosea* L., Malvaceae, Perenn., Meso-xero, Ken., Med., Ergasphig., Erg.
- Allium cepa* L., Alliaceae, Bienn., Meso, Ken., C. As. (Mayorov *et al.* 2012) or Med. (Pyšek *et al.* 2012), Ergasphig., Erg.
- Allium fistulosum* L., Alliaceae, Perenn., Meso, Ken., E. As., Ergasphig., Erg.
- Allium porrum* L., Alliaceae, Ann. or bienn., Meso, Ken., E. Med., Ergasphig., Erg.
- Allium sativum* L., Alliaceae, Bienn., Meso, Ken., Mid. As., Ergasphig., Erg.
- Allium victorialis* L., Alliaceae, Perenn., Xero-meso, Ken. (Crimea)/natural species in Ukraine (Carpathian), sub.-Alp., Col., Erg.
- Althaea officinalis* L., Malvaceae, Perenn., Hygro-meso, Arch., Iran.-Tur., Agr.-epoec., Erg.
- Amaranthus caudatus* L., Amaranthaceae, Ann., Xero-meso, Ken., S. Am. (tropic.), Ergasphig., Erg.
- Amaranthus caudatus* L. ssp. *saueri* Jehlik, Amaranthaceae, Ann., Xero-meso, Ken., S. Am., Ergasphig., Erg.
- Amaranthus cruentus* L., Amaranthaceae, Ann., Xero-meso, Ken., Trop. Am. S. Am. (tropic.), Epoec., Erg.
- Amaranthus hypochondriacus* L., Amaranthaceae, Ann., Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Amaranthus spinosus* L., Amaranthaceae, Ann., Xero-meso, Ken., S. Am., Ephem., Erg.
- Amberboa moschata* (L.) DC., Asteraceae, Ann., Xero, Ken., W. As., Ergasphig., Erg.
- Ambrosia artemisiifolia* L., Asteraceae, Ann., Xero-meso, Ken. (1925), N. Am., Agr.-epoec., Erg.-kseno; invasive; transformer (forest-steppe and steppe regions of Ukraine).
- Ambrosia trifida* L., Asteraceae, Ann., Xero-meso, Ken., N. Am., Ephem., Erg.
- Amelanchier canadensis* (L.) Medik., Rosaceae, Shrub or tree, Meso, Ken., N. Am., Ephem., Erg.
- Amelanchier ovalis* Medik., Rosaceae, Shrub, Meso, Ken., Med., Ergasphig., Erg.
- Amelanchier spicata* (Lam.) K. Koch, Rosaceae, Shrub, Meso, Ken., N. Am., Agr., Erg.; invasive.
- Ammi majus* L., Apiaceae, Bienn., Meso-xero, Ken., Med., Ephem., Erg.
- Amorpha fruticosa* L., Fabaceae, Shrub or tree, Meso, Ken., N. Am., Agr.-epoec., Erg.; invasive; transformer (central regions of Ukraine).
- Anacyclus clavatus* (Desf.) Pers., Asteraceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Anacyclus officinarum* Hayne, Asteraceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Anemone fasciculata* L., Ranunculaceae, Perenn., Meso, Ken. (Crimea – 1981), Cauc., Col., Erg.
- Anethum graveolens* L., Apiaceae, Ann., Meso, Ken., Med.-Iran.-Tur., Ergasphig., Erg.
- Anredera cordifolia* (Ten.) Steenis, Basellaceae, Liane, Meso, Ken., C. and S. Am., Ergasphig., Erg.
- Anthericum liliago* L., Asphodelaceae, Perenn., Xero-meso, Ken., Med., Ergasphig., Erg.
- Antirrhinum majus* L., Scrophulariaceae, Ann., Xero-meso, Ken., Med., Ergasphig. (Ukraine)/Agr. (Crimea), Erg.
- Apium graveolens* L., Apiaceae, Ann. or bienn., Meso, Ken., W. Eur. (Protopopova 1991) or unknown (Pyšek *et al.* 2012), Ergasphig., Erg.
- Apocynum cannabinum* L., Apocynaceae, Perenn., Meso, Ken. (2000), N. Am., Ergasphig., Erg.
- Aquilegia vulgaris* L., Ranunculaceae, Perenn., Xero-meso, Ken. (1898), W. Eur., Agr., Erg.
- Armeniaca vulgaris* Lam., Rosaceae, Tree, Meso-xero, Ken., E. and C. As., Ephem., Erg.
- Armoracia rusticana* (Lam.) Gaertn., Mey. & Scherb., Brassicaceae, Perenn., Xero-meso, Ken., Iran.-Tur., Col., Erg.
- Aronia melanocarpa* (Michx.) Elliot, Rosaceae, Shrub, Meso, Ken., N. Am., Ergasphig., Erg.
- Aronia × mitschurinii* A. K. Skvortsov & Yu. K. Maitulina, Rosaceae, Shrub, Meso, Ken., hybrid origin, Ergasphig., Erg.
- Aronia × prunifolia* (Marshall) Rehder, Rosaceae, Shrub, Meso, Ken., hybrid origin, Ergasphig., Erg.
- Arrhenatherum elatius* (L.) J. Presl & C. Presl, Poaceae, Perenn., Meso, Ken., W. Eur., Agr.-epoec., Erg.; invasive.
- Artemisia abrotanum* L., Asteraceae, S.-shrub, Xero-meso, Arch., E. Med., Agr., Erg.-kseno.
- Artemisia dracunculus* L., Asteraceae, Perenn., Meso, Ken., As., Col., Erg.
- Asclepias syriaca* L., Asclepiadaceae, Perenn., Xero-meso, Ken. (1887), N. Am., Epoec., Erg., invasive (central regions of Ukraine).
- Astrantia maxima* Pall., Apiaceae, Peren., Meso, Ken. (Crimea), Cauc.-Eur. (Bagrikova 2013), Col., Erg.
- Atriplex hortensis* L., Chenopodiaceae, Ann., Meso, Ken., As., Col., Erg.
- Avena sativa* L., Poaceae, Ann., Meso, Ken., S. Eur., Ephem., Erg.
- Balsamita major* Desf., Asteraceae, Perenn., Meso, Ken., As. (Tzvelev 1994) or Eur.-Med. (Pyšek *et al.* 2012), Col., Erg.
- Berberis amurensis* Rupr. ex Maxim., Berberidaceae, Shrub, Meso, Ken., E. As., Ergasphig., Erg.
- Berberis thunbergii* DC., Berberidaceae, Shrub, Meso, Ken., E. As., Ergasphig., Erg.
- Berberis vulgaris* L., Berberidaceae, Shrub, Meso, Ken., E. As., Col., Erg.
- Beta vulgaris* L., Chenopodiaceae, Ann. or bienn., Meso, Ken., Med. or hybrid (Pyšek *et al.* 2012), Ephem., Erg.
- Borago officinalis* L., Boraginaceae, Ann., Meso, Ken., Med., Col., Erg.
- Brassica juncea* (L.) Czern., Brassicaceae, Ann., Meso-xero, Ken., S.-E. As., Ergasphig., Erg.

- Brassica napus* L., Brassicaceae, Ann. or bienn., Xero-meso, Ken., S. Eur., Ergasphig., Erg.
- Brassica nigra* (L.) W. D. J. Koch, Brassicaceae, Ann., Xero-meso, Ken., Med., Epoec., Erg.
- Brassica oleracea* L., Brassicaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Brassica rapa* L., Brassicaceae, Ann., Xero-meso, Ken., W. As., Ergasphig., Erg.
- Broussonetia papyrifera* (L.) Vent., Moraceae, Tree or shrub, Xero-meso, Ken., E. As., Ergasphig., Erg.
- Brunnera macrophylla* (Adams) I. M. Johnst., Boraginaceae, Perenn., Xero-meso, Ken., Eur.-Med. (Pyšek *et al.* 2012), Ergasphig., Erg.
- Buddleja davidii* Franch., Buddlejaceae, Shrub, Meso, Ken. (Crimea), As., Col., Erg.
- Bupleurum fruticosum* L., Apiaceae, Shrub, Meso-xero, Ken., Med., Agr., Erg., invasive; transformer (Crimea).
- Calendula officinalis* L., Asteraceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Callistephus chinensis* (L.) Nees, Asteraceae, Ann. or bienn., Meso, Ken., E. As., Ergasphig., Erg.
- Camelina sativa* (L.) Crantz ssp. *sativa*, Brassicaceae, Ann., Xero-meso, Arch., unknown (Pyšek *et al.* 2012), Epoec., Erg.
- Campanula medium* L., Campanulaceae, Bienn., Meso, Ken., S. Eur., Ergasphig., Erg.
- Campsip grandiflora* (Thunb.) K. Schumann, Bignoniaceae, Liana, Meso, Ken., E. As., Ergasphig., Erg.
- Campsip radicans* (L.) Seem., Bignoniaceae, Liana, Meso, Ken., N.Am., Ergasphig., Erg.
- Cannabis sativa* L., Cannabaceae, Ann., Xero-meso, Ken., E. As., Epoec., Erg.
- Caragana arborescens* Lam., Fabaceae, Tree or shrub, Meso, Ken., Sib., Ergasphig., Erg.
- Carthamus tinctorius* L., Asteraceae, Ann., Meso-xero, Ken., S. As. (Mayorov *et al.* 2012), Ergasphig., Erg.
- Castanea sativa* Mill., Fagaceae, Tree, Meso, Ken., Cauc.-Asia Minor-Med., Ergasphig., Erg.
- Catalpa bignonioides* Walt., Bignoniaceae, Tree, Meso, Ken., N. Am., Ergasphig., Erg.
- Cedrus atlantica* (Endl.) Carrière, Pinaceae, Tree, Xero-meso, Ken. (Crimea – 1995), N.-W. Afr., Ergasphig., Erg.
- Cedrus deodara* (Roxb.) G. Don, Pinaceae, Tree, Xero-meso, Ken. (Crimea), C. As., Col., Erg.
- Celtis australis* L., Celtidaceae, Tree, Xero-meso, Ken. (Crimea), Med., Ergasphig., Erg.
- Celtis caucasica* Willd., Celtidaceae, Tree, Xero-meso, Ken. (Crimea), Cauc., Ergasphig., Erg.
- Cenchrus longispinus* (Hack.) Fernald, Poaceae, Ann., Meso-xero, Ken. (1950), N. Am., Agr.-epoec., Erg.-kseno; invasive.
- Centranthus ruber* (L.) DC., Valerianaceae, Perenn., Xero-meso, Ken. (Crimea), Med., Agr., Erg.
- Cerastium tomentosum* L., Caryophyllaceae, Perenn., Meso-xero, Ken. (Crimea), Med., Col., Erg.
- Cerasus besseyi* (Bailey) Sok., Rosaceae, Shrub, Meso, Ken., N. Am., Ergasphig., Erg.
- Cerasus tomentosa* (Thunb.) Wall., Rosaceae, Tree or shrub, Meso, Ken., E. As., Ergasphig., Erg.
- Cerasus vulgaris* Mill., Rosaceae, Tree or shrub, Meso, Ken., E. Med. or Balc., Ergasphig., Erg.
- Ceratochloa carinata* (Hook. & Arn.) Tutin, Poaceae, Bienn., Xero-meso, Ken., N. Am., Ephem., Erg.
- Ceratostigma plumbaginoides* Bunge, Plumbaginaceae, Perenn., Xero-meso, Ken., As., Col., Erg.
- Cercis siliquastrum* L., Caesalpiniaceae, Shrub, Xero-meso, Ken. (1809), Med., Ergasphig., Erg.
- Chaenomeles japonica* (Thunb.) Lindl., Rosaceae, Shrub, Meso, Ken., E. As., Ergasphig., Erg.
- Chamaemelum nobile* (L.) All., Asteraceae, Perenn., Meso, Ken., Med., Ergasphig., Erg.
- Cheiranthus cheiri* L., Brassicaceae, Perenn., Xero-meso, Ken., Med., Ergasphig., Erg.
- Chenopodium capitatum* (L.) Ambrosi, Chenopodiaceae, Ann., Xero-meso, Ken., W. Eur., Ergasphig., Erg.
- Chenopodium foliosum* Asch., Chenopodiaceae, Ann., Xero-meso, Ken., Eur.-Med.-As., Epoec., Erg.
- Chenopodium schraderianum* Schult., Chenopodiaceae, Ann., Meso, Ken., Afr. (tropic and subtropic), Ergasphig., Erg.
- Chrysanthemum carinatum* Schousb., Asteraceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Chrysanthemum coronarium* L., Asteraceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Cicer arietinum* L., Fabaceae, Ann., Ken., Meso, Med., Ephem., Erg.
- Citrullus lanatus* (Thunb.) Matsum. & Nakai, Cucurbitaceae, Ann., Xero-meso, Ken., Afr., Ergasphig., Erg.
- Cleome spinosa* Jacq., Cleomaceae, Ann., Xero-meso, Ken. (2013), S. Am., Ergasphig., Erg.
- Clematis flammula* L., Ranunculaceae, Liana, Xero-meso, Ken., Med., Col., Erg.
- Clematis vitalba* L., Ranunculaceae, Liana, Xero-meso, Ken., S. or C. Eur., Ergasphig., Erg.
- Cnicus benedictus* L., Asteraceae, Ann., Meso-xero, Ken., S. Eur., Ergasphig., Erg.
- Colutea arborescens* L., Fabaceae, Shrub, Meso-xero, Ken. (Crimea)/natural species in Ukraine, Eur.-Med. (Pyšek *et al.* 2012), Col., Erg.
- Colutea orientalis* Mill., Fabaceae, Tree, Xero, Ken. (Crimea), unknown, Agr., Erg.
- Commelinia communis* L., Commelinaceae, Ann., Meso, Ken. (1946), S.-E. As., Ephem., Erg.-kseno.
- Coreopsis grandiflora* Hogg ex Sweet, Asteraceae, Ann. or bienn., Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Coreopsis tinctoria* Nutt., Asteraceae, Ann. or bienn., Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Coriandrum sativum* L., Apiaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Cosmos bipinnatus* Cav., Asteraceae, Ann., Xero-meso, Ken., N. Am., Col., Erg.
- Cotinus coggygria* Scop., Anacardiaceae, Tree or shrub, Xero, Ken. (steppe regions of Ukraine)/natural species in Crimea, Med., Ergasphig., Erg.
- Crocus flavus* Weston, Iridaceae, Perenn., Xero-meso, Ken., unknown (Pyšek *et al.* 2012), Erg.-lip., Erg.
- Cucumis sativus* L., Cucurbitaceae, Ann., Meso, Ken., S.-E. As., Ergasphig., Erg.
- Cucurbita maxima* Duch., Cucurbitaceae, Ann., Meso, Ken., S. Am., Ergasphig., Erg.
- Cucurbita pepo* L., Cucurbitaceae, Ann., Meso, Ken., N. Am., Ergasphig., Erg.
- Cupressus sempervirens* L., Cupressaceae, Tree, Meso-xero, Ken. (Crimea – 1995), Med., Col., Erg.
- Cyclamen hederifolium* Ait., Primulaceae, Perenn., Meso, Ken., Med., Col., Erg.
- Cyclamen vernum* Sweet, Primulaceae, Perenn., Meso, Ken., Med., Ergasphig., Erg.
- Cydonia oblonga* Mill., Rosaceae, Shrub, Xero-meso, Ken. (Crimea), W. As., Col., Erg.
- Cymbalaria muralis* Gaertn., Mey. & Scherb., Scrophulariaceae, Bienn., Meso-xero, Ken., Med., Col., Erg.
- Daphne laureola* L., Thymelaeaceae, Shrub, Meso, Ken., Med., Ergasphig., Erg.
- Datura innoxia* Mill., Solanaceae, Ann., Xero-meso, Ken., N. Am., Ephem., Erg.
- Datura meteloides* DC., Solanaceae, Ann., Xero-meso, Ken., N. Am., Ergasphig., Erg.

- Datura tatula* L., Solanaceae, Ann., Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Dianthus barbatus* L., Caryophyllaceae, Perenn., Meso, Ken., Middle Eur., Ergasphig., Erg. Рогович
- Digitalis lanata* Ehrh., Scrophulariaceae, Perenn., Meso, Ken. (2001), S. Eur.-Balcan, Col., Erg.
- Digitalis purpurea* L., Scrophulariaceae, Ann., Ken., Meso, Eur.-Med., Ergasphig., Erg.
- Diospyros lotus* L., Ebenaceae, Tree, Xero-meso, Ken. (Crimea), E. As., Col., Erg.
- Dipsacus sativus* (L.) Honck., Dipsacaceae, Bienn., Xero-meso, Ken., unknown (Pyšek *et al.* 2012), Ergasphig., Erg.
- Disphania schraderiana* (Schult.) Mosyakin & Clemants, Chenopodiaceae, Ann., Xero-meso, Ken., Afr. (tropic and subtropic), Ergasphig., Erg.
- Dracocephalum moldavica* L., Lamiaceae, Ann., Xero-meso, Ken., E. As., Ergasphig., Erg.
- Duchesnea indica* (Andr.) Focke, Rosaceae, Perenn., Xero-meso, Ken., S.-E. As., Ergasphig., Erg.
- Echinacea purpurea* (L.) Moench, Asteraceae, Perenn., Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Echinochloa esculenta* (A. Br.) H. Scholz, Poaceae, Ann., Xero-meso, Ken., E. As., Ergasphig., Erg.
- Echinocystis lobata* (Michx.) Torr. & Gray, Cucurbitaceae, Ann., Meso, Ken. (1929), N. Am., Agr.-epoec., Erg.; invasive; transformer (Transcarpathia).
- Elaeagnus angustifolia* L., Elaeagnaceae, Tree or shrub, Xero-meso, Ken., Med., Agr., Erg.; invasive; transformer (Black Sea Region).
- Elaeagnus commutata* Bernh. ex Rydb., Elaeagnaceae, Tree or shrub, Meso, Ken., N. Am., Col., Erg.
- Elodea canadensis* Michx., Hydrocharitaceae, Perenn., Hydro, Ken. (1894), N. Am., Agr., Erg., invasive.
- Elodea densa* (Planch.) Caspary, Hydrocharitaceae, Perenn., Hydro, Ken. (2001), S. Am., Ephem., Erg.
- Elodea nuttallii* (Planchon) H. St. Joh. Caspary, Hydrocharitaceae, Perenn., Hydro, Ken. (2004), N. Am., Ephem., Erg.
- Eruca vesicaria* (L.) Cav., Brassicaceae, Ann., Xero-meso, Ken., E. Med., Ephem., Erg.
- Eschscholzia californica* Cham., Papaveraceae, Ann., Meso-xero, Ken., N. Am., Ergasphig., Erg.
- Eudianthe coeli-rosa* (L.) Rchb., Caryophyllaceae, Ann., Xero-mezo, Ken. (Crimea), W. Med., Ephem., Kseno/Erg.
- Euonymus japonica* Thunb., Celastraceae, Shrub, Meso, Ken., E. As., Ergasphig., Erg.
- Euonymus sacrosancta* Koidz., Celastraceae, Shrub, Meso, Ken., E. As., Ergasphig., Erg.
- Euphorbia lathyris* L., Euphorbiaceae, Ann., Meso-xero, Ken., Med., Ergasphig., Erg.
- Euphorbia marginata* Pursh, Euphorbiaceae, Ann., Meso-xero, Ken., N. Am., Ergasphig., Erg.
- Fagopyrum esculentum* Moench, Polygonaceae, Ann., Meso, Ken., C. As., Ergasphig., Erg.
- Fallopia baldschuanica* (Regel) Holub, Polygonaceae, Liana, Xero-meso, Ken., Middle As., Ergasphig., Erg.
- Ficus carica* L., Moraceae, Tree or shrub, Xero, Ken., As.-Med., Epoec., Erg.
- Flueggea suffruticosa* (Pall.) Baillon, Euphorbiaceae, Shrub, Meso-xero, Ken. (1999), E. As., Col., Erg.
- Foeniculum vulgare* Mill., Apiaceae, Bienn., Xero-meso, Ken., Med., Col., Erg.
- Fragaria ×ananassa* (Duchesne) Duchesne, Rosaceae, Perenn., Meso, Ken., hybrid origin, Ergasphig., Erg.
- Fraxinus ornus* L., Oleaceae, Tree, Meso-xero, Ken., Med., Agr. Erg.; invasive.
- Fraxinus pennsylvanica* Marsh., Oleaceae, Oleaceae, Tree, Xero-meso, Ken., N. Am., Col., Erg.
- Fritillaria ophioglossifolia* Freun. & Sint., Liliaceae, Perenn., Meso, Ken. (Crimea), Cauc., Col., Erg.
- Gaillardia pulchella* Foug., Asteraceae, Ann. or bienn., Meso-xero, Ken., Am., Col., Erg.
- Gleditsia triacanthos* L., Caesalpiniaceae, Tree, Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Glycine max* (L.) Merr., Fabaceae, Ann., Ken., Xero-meso, E. As., Ergasphig., Erg.
- Grossularia uva-crispa* (L.) Mill. ssp. *reclinata* (L.) Dostal, Grossulariaceae, Shrub, Meso, Ken., Eur., Ergasphig., Erg.
- Gypsophila acutifolia* Fisch. ex Spreng., Caryophyllaceae, Perenn., Xero-meso, Ken., Cauc., Ergasphig., Erg.
- Gypsophila elegans* M. Bieb., Caryophyllaceae, Ann., Xero-meso, Ken., Cauc.-Asia Minor or Eur.-Med. (Pyšek *et al.* 2012), Ephem., Erg.
- Gypsophila paulii* Klok., Caryophyllaceae, Perenn., Xero-meso, Ken. (S.-E. Ukraine, Ostapko, Boyko, Mosyakin 2010), S. Cis Black Sea, Ergasphig., Erg.
- Halimodendron halodendron* (Pall.) Voss, Fabaceae, Shrub, Meso, Ken., unknown, Ephem., Erg.
- Helianthus annuus* L. var. *macrocarpa* Lucznik, Asteraceae, Ann., Xero-meso, Ken., N. Am., Ephem., Erg.
- Helianthus decapetalus* L., Asteraceae, Perenn., Meso, Ken., N. Am., Epoec., Erg.
- Helianthus ×laetiflorus* Pers., Asteraceae, Perenn., Xero-meso, Ken., N. Am., Agr.-epoec., Erg.; invasive.
- Helianthus rigidus* (Cass.) Desf., Asteraceae, Perenn., Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Helianthus strumosus* L., Asteraceae, Perenn., Meso, Ken., N. Am., Col., Erg.
- Helianthus subcanescens* (A.Gray) E. E. Wats., Asteraceae, Perenn., Meso, Ken., N. Am., Agr.-epoec., Erg.; invasive.
- Helianthus tuberosus* L., Asteraceae, Perenn., Meso, Ken., N. Am., Agr.-epoec., Erg.; invasive; transformer (Transcarpathia).
- Helichrysum italicum* (Roth) G. Don fil., Asteraceae, S.-shrub, Xero, Ken. (Crimea – 1995), Med., Col., Erg.
- Heliopsis scabra* Dun., Asteraceae, Perenn., Xero-meso, Ken., N. Am., Col., Erg.
- Helleborus dumetorum* Waldst. & Kit., Ranunculaceae, Perenn., Meso, Ken., S. Eur., Ergasphig., Erg.
- Helleborus niger* L., Ranunculaceae, Perenn., Meso, Ken., S. Eur., Ergasphig., Erg. Note. The species included in to Red Data Book of Ukraine, according to S. Mosyakin and M. Fedorochuk (1999) probably cultivated and escaped.
- Helleborus viridis* L., Ranunculaceae, Perenn., Meso, Ken., S. Eur., Ergasphig., Erg.
- Hemerocallis fulva* (L.) L., Hemerocallidaceae, Perenn., Xero-meso, Ken., E. As., Ergasphig., Erg.
- Heracleum mantegazzianum* Sommier & Levier, Apiaceae, Perenn., Hygro, Ken. (1964), Cauc., Agr.-epoec., Erg.; invasive; transformer (Carpathian).
- Heracleum sosnowskyi* Manden., Apiaceae, Perenn., Hygro, Ken., Cauc., Agr.-epoec., Erg., invasive; transformer (Carpathian, Polissya [Forest zone of Ukraine]).
- Hesperis matronalis* L., Brassicaceae, Ann. or bienn., Xero-meso, Ken. (left forest and steppe regions of Ukraine), Sub.-Med. (Ilyinska *et al.* 2007) or Eur. and Med. (Pyšek *et al.* 2012), Ergasphig., Erg.
- Hesperis pycnотricha* Borbas & Degen, Brassicaceae, Bienn., Meso, Ken., Cauc.-As. min., Ergasphig., Erg.
- Hesperis sibirica* L., Brassicaceae, Bienn., Meso, Ken., As., Ergasphig., Erg.
- Hordeum distichon* L., Poaceae, Ann., Xero-meso, Ken., E. Med., Ergasphig., Erg.

- Hordeum jubatum* L., Poaceae, Ann., Meso-xero, Ken., N. Am., Ephem., Erg.
- Hordeum vulgare* L., Poaceae, Ann., Xero-meso, Ken., As., Ergasphig., Erg.
- Humulus japonicus* Siebold & Zucc., Cannabaceae, Liana, Meso, Ken. (2002), E. As., Ergasphig., Erg.
- Hypericum calycinum* L., Hypericaceae, Shrub, Xero-meso, Ken., E. Med., Ergasphig., Erg.
- Hyssopus officinalis* L., Lamiaceae, Perenn., Xero-meso, Ken., Med., Ergasphig., Erg.
- Iberis amara* L., Brassicaceae, Ann., Xero-meso, Ken., Med. (Protopopova, 1991), Ergasphig., Erg.
- Iberis pinnata* L., Brassicaceae, Ann., Xero, Ken., W. Med., Ergasphig., Erg.
- Iberis umbellata* L., Brassicaceae, Ann. or bienn., Xero-meso, Ken., Med., Ergasphig., Erg.
- Impatiens balfourii* Hook. f., Balsaminaceae, Ann., Meso, Ken. (2013), W. Hymal., Ephem., Erg.
- Impatiens glandulifera* Royle, Balsaminaceae, Ann., Hydro-meso, Ken. (1939), S.-E. As., Agr.-epoec., Erg.; invasive; transformer (Transcarpathia).
- Ipomoea purpurea* (L.) Roth, Convolvulaceae, Ann., Xero-meso, Ken., Trop. Am., Ergasphig., Erg.
- Iris ×germanica* L., Iridaceae, Perenn., Meso, Ken. (Crimea), hybrid or Eur.-As. (Pyšek *et al.* 2012), Col., Erg.
- Iris musulmanica* Fomin, Iridaceae, Perenn., Meso, Ken., Cauc. or Eur. (Bagrikova 2013), Ergasphig., Erg.
- Iris pallida* Lam., Iridaceae, Perenn., Meso, Ken., Med., Ergasphig., Erg.
- Iva xanthiifolia* Nutt., Asteraceae, Perenn., Xero-meso, Ken. (1842), N. Am., Epoec., Erg.; invasive.
- Jasminum nudiflorum* Lindl., Oleaceae, Shrub, Xero-meso, Ken., E. As., Ergasphig., Erg.
- Jasminum officinale* L., Oleaceae, Shrub, Meso, Ken., E. Med., Ergasphig., Erg.
- Juglans cinerea* L., Juglandaceae, Tree, Meso, Ken., N. Am., Ergasphig., Erg.
- Juglans mandshurica* Maxim., Juglandaceae, Tree, Meso, Ken., As., Ergasphig., Erg.
- Juglans nigra* L., Juglandaceae, Tree, Meso, Ken., N. Am., Ergasphig., Erg.
- Juglans regia* L., Juglandaceae, Tree, Meso, Ken., Balcan-C. As., Ergasphig., Erg.
- Kali collina* (Pall.) Akhani & E. H. Roalson, Chenopodiaceae, Ann., Xero, Ken., Tur., Ergasphig., Erg.
- Kochia scoparia* (L.) Schrad., Chenopodiaceae, Ann., Meso-xero, Ken., Iran.-Tur., Ergasphig., Erg.
- Koelreuteria paniculata* Laxm., Sapindaceae, Tree or shrub, Meso, Ken. (Crimea), E. As., Col., Erg.
- Laburnum anagyroides* Medik., Fabaceae, Trees or shrub, Xero-meso, Ken., S. Eur., Ergasphig., Erg.
- Lactuca sativa* L., Asteraceae, Bienn., Meso, Ken., unknown (Pyšek *et al.* 2012), Ergasphig., Erg.
- Lagurus ovatus* L., Poaceae, Ann., Xero, Ken. (Crimea), Med. (Pyšek *et al.* 2012), Ephem., Kseno./Erg.
- Lallemandia iberica* (M. Bieb.) Fisch. & C. A. Mey., Lamiaceae, Ann., Meso-xero, Ken., W. As., Ergasphig., Erg.
- Lamprocapnos spectabilis* (L.) Fukuhara (Fumariaceae), Perenn., Meso, Ken., E. As. (Mayorov *et al.* 2012), Ergasphig., Erg.
- Lathyrus odoratus* L., Fabaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Lathyrus sativus* L., Fabaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Laurocerasus officinalis* M. Roem., Rosaceae, Tree or shrub, Xero-meso, Ken., Med.-Iran.-Tur., Agr., Erg.
- Laurus nobilis* L., Lauraceae, Tree or shrub, Xero-meso, Ken., Med., Ergasphig., Erg.
- Lavandula angustifolia* Mill., Lamiaceae, Perenn., Xero-meso, Ken., Med. (Pyšek *et al.* 2012), Ergasphig., Erg.
- Lavatera trimestris* L., Malvaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Lens culinaris* Medik., Fabaceae, Ann., Meso, Ken., unknown (Pyšek *et al.* 2012), Ergasphig., Erg.
- Lepidium sativum* L., Brassicaceae, Ann., Meso, Ken., Med., Ergasphig., Erg.
- Levisticum officinale* W. D. J. Koch, Apiaceae, Perenn., Xero-meso, Ken., E. Med., Ergasphig., Erg.
- Leucanthemum maximum* (Ramond) DC., Asteraceae, Perenn., Xero-meso, Ken., W. Eur., Ergasphig., Erg.
- Lilium monadelphum* M. Bieb., Liliaceae, Perenn., Xero-meso, Ken. (Crimea), Cauc., Ephem., Erg.
- Linaria bipartita* (Vent.) Willd., Scrophulariaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Linaria incarnata* (Vent.) Spreng., Scrophulariaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Linum grandiflorum* Desf., Linaceae, Ann., Meso-xero, Ken., N. Afr., Col., Erg.
- Linum usitatissimum* L., Linaceae, Ann., Meso-xero, Ken., unknown, Ephem., Erg.
- Lobularia maritima* (L.) Desv., Brassicaceae, Ann., Meso-xero, Ken., Med., Ergasphig., Erg.
- Lolium multiflorum* Lam., Poaceae, Ann. or bienn., Meso, Ken., Med.-Iran.-Tur., Epoec., Erg.
- Lonicera caprifolium* L., Caprifoliaceae, Liana, Xero-meso, Ken., Med., Ergasphig., Erg.
- Lonicera etrusca*, Caprifoliaceae, Shrub, Xero-meso, Ken., Med., Agr. (Crimea), Erg.
- Lonicera standishii* Jacq., Caprifoliaceae, Shrub, Xero-meso, Ken. (Crimea), As., Col., Erg.
- Lonicera tatarica* L., Caprifoliaceae, Shrub or tree, Meso, Ken., As., Ergasphig., Erg.
- Lunaria annua* L., Brassicaceae, Ann., Xero-meso, Ken., S. Eur., Ergasphig., Erg.
- Lupinus luteus* L., Fabaceae, Ann., Meso, Ken., W. Med., Ergasphig., Erg.
- Lupinus perennis* L., Fabaceae, Perenn., Meso, Ken. N. Am., Ephem., Erg.
- Lupinus polyphyllus* Lindl., Fabaceae, Perenn., Meso, Ken., N. Am., Agr.-epoec., Erg.; invasive.
- Lychnis chalcedonica* L., Caryophyllaceae, Perenn., Meso, Ken., As., Ergasphig., Erg.
- Lycium barbatum* L., Solanaceae, Shrub, Meso-xero, Arch., E. As., Epoec., Erg.
- Lycopersicon esculentum* Mill. s.l., Solanaceae, Ann., Meso, Ken., S. Am., Ergasphig., Erg.
- Maclura pomifera* (Rafin.) Schneid., Moraceae, Tree, Meso, Ken. (Crimea), N. Am., Col., Erg.
- Mahonia aquifolium* (Pursh) Nutt., Berberidaceae, Shrub, Xero-meso, Ken., N. Am., Agr. (Polissya [Forest zone of Ukraine], Crimea), Erg.
- Malope trifida* Cav., Malvaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Malus domestica* Borkh., Rosaceae, Tree, Meso, Ken., unknown, Ergasphig., Erg.
- Malva mauritiana* L., Malvaceae, Bienn., Xero-meso, Ken., Med., Epoec., Erg.
- Malva moschata* L., Malvaceae, Perenn., Xero-meso, Ken., Med. or As. min. (Protopopova 1991), Ergasphig., Erg.
- Malva verticillata* L., Malvaceae, Ann., Xero-meso, Ken., As., Ephem., Erg.
- Matthiola bicornis* (Sibth. & Smith) DC., Brassicaceae, Ann. or bienn., Xero-meso, Ken., Med., Ergasphig., Erg.

- Matthiola incana* (L.) R. Br., Brassicaceae, Ann. or bienn., Meso, Ken., S. Eur., Ergasphig., Erg.
- Matthiola longipetala* (Vent.) DC., Brassicaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Medicago sativa* L., Fabaceae, Perenn., Xero-meso, Ken., E. Med., Epoec., Erg.
- Melissa officinalis* L., Lamiaceae, Perenn., Xero-meso, Ken., Med., Ergasphig., Erg.
- Melo sativus* Sager. ex M. Roem., Cucurbitaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Mentha ×gracilis* Sole, Lamiaceae, Perenn., Meso, Ken., hybrid origin, Ephem., Erg.
- Mentha ×piperita* L., Lamiaceae, Perenn., Meso, Ken., hybrid origin, Ephem., Erg.
- Mentha spicata* L., Lamiaceae, Perenn., Hygro, Ken., Med., Ergasphig., Erg.
- Mimulus guttatus* DC., Scrophulariaceae, Perenn., Meso, Ken., N. Am., Ergasphig., Erg.
- Misanthus sinensis* Anderson, Poaceae, Perenn., Xero-meso, Ken., E. As., Ergasphig., Erg.
- Molucella laevis* L., Lamiaceae, Ann., Xero-meso, Ken., Med.-Iran.-Tur., Ergasphig., Erg.
- Morus alba* L., Moraceae, Tree, Xero-meso, Ken., E. As., Ergasphig., Erg.
- Morus nigra* L., Moraceae, Tree, Xero-meso, Ken., As., Ergasphig., Erg.
- Morus rubra* L., Moraceae, Tree, Xero-meso, Ken. (Crimea), N. Am., Col., Erg.
- Myrrhis odorata* (L.) Scop., Apiaceae, Perenn., Meso, Ken., W. Eur., Ergasphig., Erg.
- Narcissus poëticus* L., Amaryllidaceae, Perenn., Xero-meso, Ken., Med. (Pyšek *et al.* 2012) or Eur. (Mayorov *et al.* 2012), Ergasphig., Erg.
- Narcissus pseudonarcissus* L., Amaryllidaceae, Perenn., Xero-meso, Ken., W. Eur., Ergasphig., Erg.
- Nicandra physalodes* (L.) Gaertn., Solanaceae, Ann., Xero-meso, Ken. (1855), S. Am., Ergasphig., Erg.
- Nicotiana alata* Link & Otto, Solanaceae, Ann., Meso, Ken., S. Am., Ergasphig., Erg.
- Nicotiana tabacum* L., Solanaceae, Ann., Meso, Ken, Trop. Am., Ergasphig., Erg.
- Nicotiana rustica* L., Solanaceae, Ann., Meso, Ken (1869), N. Am., Ergasphig., Erg.
- Nigella sativa* L., Ranunculaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Oenothera glazioviana* Michelii, Onagraceae, Bienn., Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Oenothera suaveolens* Desf., Onagraceae, Bienn., Xero-meso, Ken., unknown, Ergasphig., Erg.
- Olea europaea* L., Oleaceae, Tree or shrub, Meso-xero, Ken., E. Med., Ergasphig., Erg.
- Omphalodes linifolia* (L.) Moench, Boraginaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Omphalodes verna* Moench, Boraginaceae, Perenn., Xero-meso, Ken., Med., Epoec., Erg.
- Onobrychis viciifolia* Scop., Fabaceae, Perenn., Xero-meso, Ken., S. Eur., Ephem., Erg.
- Opuntia fragilis* (Nutt.) Haw., Cactaceae, Perenn., Xero, Ken. (Crimea), C. Am., Agr., Erg.
- Opuntia humifusa* (RAfr.) RAfr., Cactaceae, Perenn., Xero, Ken. (Crimea – 2004), N. Am., Ergasphig., Erg.
- Opuntia lindheimeri* Engel., Cactaceae, Perenn., Xero, Ken. (Crimea), N. Am., Agr., Erg.
- Opuntia phaeocantha* Engelm., Cactaceae, Perenn., Xero, Ken. (Crimea – 2004), N. Am., Ergasphig., Erg.
- Opuntia tortispina* Engelm., Cactaceae, Perenn., Xero, Ken. (Crimea – 2004), N. Am., Ergasphig., Erg.
- Opuntia vulgaris* Mill., Cactaceae, Perenn., Xero, Ken. (Crimea – 2004), S. Am., Agr., Erg.
- Ornithogalum kochii* Parl., Hyacinthaceae, Perenn., Xero, Ken. (forest and forest-steppe regions of Ukraine), N. Pann., Ergasphig., Erg.
- Ornithopus sativus* Brot., Fabaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Padus serotina* (Ehrh.) Ag., Rosaceae, Shrub, Xero-meso, Ken., N. Am., Agr., Erg.
- Padus virginiana* (L.) Roem., Rosaceae, Tree, Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Panicum miliaceum* L., Poaceae, Ann., Xero-meso, Ken., S.-E. As., Ergasphig., Erg.
- Papaver somniferum* L., Papaveraceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Parthenocissus inserta* (A. Kern.) Fritsch, Vitaceae, Liana, Meso, Ken., N. Am., Agr.-Epoec., Erg.
- Parthenocissus quinquefolia* (L.) Planch., Vitaceae, Liana, Meso, Ken., N. Am., Col., Erg.
- Pastinaca sativa* L., Apiaceae, Bienn., Meso, Ken., Eur.-As., Ephem., Erg.
- Persicaria orientalis* (L.) Spach, Polygonaceae, Ann., Meso, Ken., S.-E. As., Ergasphig., Erg.
- Petrohragia saxifraga* (L.) Link, Caryophyllaceae, Ann., Xero, Ken., Med., Ergasphig., Erg.
- Petrosedum rupestre* (L.) Grulich, Crassulaceae, Perenn., Xero, Ken., Cauc., Ergasphig., Erg.
- Petroselinum crispum* (Mill.) A. W. Hill, Apiaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Petunia ×atkinsiana* D. Don ex Loudon, Solanaceae, Ann., Meso, Ken., hybrid or unknown (Pyšek *et al.* 2012), Ergasphig., Erg.
- Phacelia tanacetifolia* Benth., Hydrophyllaceae, Ann., Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Phalaris canariensis* L., Poaceae, Ann., Xero-meso, Ken., Med., Ephem., Erg.
- Phalaris minor* Retz., Poaceae, Ann., Xero-meso, Ken., Med., Ephem., Kseno./Erg.
- Phaseolus vulgaris* L., Fabaceae, Ann., Meso, Ken., S. Am., Ergasphig., Erg.
- Phellodendron amurense* Rupr., Rutaceae, Tree, Meso, Ken, E. As., Ergasphig., Erg.
- Phlox paniculata* L., Polemoniaceae, Perenn., Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Phlox subulata* L., Polemoniaceae, Perenn., Xero, Ken., N. Am., Ergasphig., Erg.
- Physalis ixocarpa* Brot. ex Hornem., Solanaceae, Ann., Meso, Ken., N. Am., Ergasphig., Erg.
- Physocarpus opulifolius* (L.) Maxim., Rosaceae, Shrub, Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Phytolacca americana* L., Phytolaccaceae, Perenn., Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Pimpinella anisum* L., Apiaceae, Ann., Xero-meso, Ken., E. Med., Ergasphig., Erg.
- Pinus banksiana* Lamb., Pinaceae, Tree, Xero, Ken., N. Am., Ergasphig., Erg.
- Pinus strobus* L., Pinaceae, Trees, Xero, Ken., N. Am., Ergasphig., Erg.
- Pistia stratiotes* L., Araceae, Perenn., Hydro, Ken., Afr., Ephem., Erg.
- Pisum arvense* L., Fabaceae, Ann., Xero-meso, Ken., unknown (Pyšek *et al.* 2012), Cauc. (Protopopova 1991), Ergasphig., Erg.
- Pisum sativum* L., Fabaceae, Ann., Meso, Ken., S. Eur., Ergasphig., Erg.
- Platanus ×hispanica* Mill. ex Muenchh., Platanaceae, Tree, Xero-meso, Ken. (Crimea), hybrid origin, Col., Erg.
- Platycladus orientalis* (L.) Franco, Cupressaceae, Tree or shrub, Xero-meso, Ken., E. As., Ergasphig., Erg.

- Polygala caucasica* Rupr., Polygalaceae, Perenn., Meso, Ken. (Crimea – 1981), Cauc., Col., Erg.
- Polygonum pensylvanicum* L., Polygonaceae, Ann., Meso, Ken. (1998), N. Am., Ergasphig., Erg.
- Populus balsamifera* L., Salicaceae, Tree, Meso, Ken., N. Am., Ergasphig., Erg.
- Populus ×canadensis* Moench, Salicaceae, Tree, Meso, Ken., hybrid origin, Ergasphig., Erg.
- Populus laurifolia* Ledeb., Salicaceae, Tree, Meso, Ken., C. As., Ergasphig., Erg.
- Populus simonii* Carr., Salicaceae, Tree, Meso, Ken., E. As., Ergasphig., Erg.
- Portulaca grandiflora* Hook., Portulacaceae, Ann., Xero-meso, Ken., S. Am., Ergasphig., Erg.
- Portulaca oleracea* L. ssp. *sativa* (Haw.) Schuebl. & G. Martens, Portulacaceae, Ann., Xero-meso, Ken., Med. (Pyšek *et al.* 2012), Ergasphig., Erg.
- Prunus cerasifera* Ehrh., Rosaceae, Tree or shrub, Xero-meso, Ken., Med., Ergasphig., Erg.
- Prunus divaricata* Ledeb., Rosaceae, Tree or shrub, Xero-meso, Ken., As., Ergasphig., Erg.
- Prunus domestica* L., Rosaceae, Tree or shrub, Xero-meso, Ken., hybrid (Tzvelev 2001), Ergasphig., Erg.
- Prunus insititia* L., Rosaceae, Shrub or tree, Xero-meso, Ken., As., Ergasphig., Erg.,
- Ptelea trifoliata* L., Rutaceae, Tree, Meso, Ken., N. Am., Ergasphig., Erg.
- Pueraria lobata* (Willd.) Ohwi, Fabaceae, Shrub, Meso, Ken. (Crimea), S.-E. As., Col., Erg.
- Pyracantha rogersiana* (A. B. Jacks.) Bean., Rosaceae, Shrub, Meso, Ken. (Crimea), As., Col., Erg.
- Pyrethrum cinerariifolium* Trev., Asteraceae, Perenn., Xero-meso, Ken., Balcan., Ergasphig., Erg.
- Pyrethrum macrophyllum* (Waldst. & Kit.) Willd., Asteraceae, Perenn., Meso, Ken., E. Med., Ergasphig., Erg.
- Pyrethrum partheniifolium* Willd., Asteraceae, Perenn., Meso-xero, Ken., Med.-Iran.-Tur., Ergasphig., Erg.
- Pyrethrum parthenium* (L.) Smith, Asteraceae, Perenn., Meso-xero, Ken., W. Eur., Ergasphig., Erg.
- Pyrus communis* L., Rosaceae, Tree, Meso, Ken., unknown (Pyšek *et al.* 2012) or As. (Bagrikova 2013), Agr., Erg./Kseno.
- Quercus ilex* L., Fagaceae, Tree, Meso-xero, Ken. (Crimea), Med., Ergasphig., Erg.
- Quercus palustris* Moench, Fagaceae, Tree, Meso, Ken., N. Am., Agr., Erg.
- Quercus rubra* L., Fagaceae, Tree, Meso, Ken., N. Am., Agr., Erg.; invasive.
- Raphanus sativus* L., Brassicaceae, Ann. or bienn., Xero-meso, Ken., Med., Ergasphig., Erg.
- Reynoutria ×bohemica* Chrtek & Chrtkova, Polygonaceae, Perenn., Meso, Ken., hybrid origin, Epoec., Erg.; invasive.
- Reynoutria japonica* Houtt., Polygonaceae, Perenn., Meso, Ken. (1929), E. As., Agr.-epoec., Erg.; invasive; transformer (Transcarpathia).
- Reynoutria sachalinensis* (Fr. Schmidt) Nakai, Polygonaceae, Perenn., Meso, Ken. (1929), E. As., Col., Erg.
- Rhamnus alaternus* L., Rhamnaceae, Shrub, Xero-meso, Ken., Med., Agr., Erg.; invasive; transformer (Crimea).
- Rhus typhina* L., Anacardiaceae, Tree, Xero, Ken., N. Am., Ergasphig., Erg.
- Ribes aureum* Pursh, Grossulariaceae, Shrub, Meso, Ken., N. Am., Ergasphig., Erg.
- Ribes rubrum* L., Grossulariaceae, Shrub, Meso, Ken., W. Eur., Ergasphig., Erg.
- Ribes spicatum* E. Robson, Grossulariaceae, Shrub, Meso, Ken. (Crimea) / natural species in Ukraine, Eur.-As. (Pyšek *et al.* 2012) or Eur. (Bagrikova 2013), Col., Erg.
- Ricinus communis* L., Euphorbiaceae, Ann., Meso, Ken., Afr., Ergasphig., Erg.
- Robinia hispida* L., Fabaceae, Shrub, Xero-meso, Ken., N. Am., Col., Erg.
- Robinia pseudoacacia* L., Fabaceae, Tree, Xero-meso, Ken., N. Am., Epoec., Erg.; invasive.
- Rosa alba* L., Rosaceae, Shrub, Xero-meso, Ken., hybrid origin, Ergasphig., Erg.
- Rosa centifolia* L., Rosaceae, Shrub, Xero-meso, Ken., hybrid origin, Ergasphig., Erg.
- Rosa multiflora* Thunb., Rosaceae, Shrub, Xero-meso, Ken., E. As., Ergasphig., Erg.
- Rosa rugosa* Thunb., Rosaceae Shrub, Xero-meso, Ken., E. As., Col., Erg.
- Rosmarinum officinalis* L., Lamiaceae, S-shrub, Meso-xero, Ken., Med., Ergasphig., Erg.
- Rubia tinctorum* L., Rubiaceae, Perenn., Meso-xero, Ken., Med.-Iran.-Tur., Col., Erg.
- Rubus idaeus* L., Rosaceae, Shrub, Meso, Ken. (Crimea)/natural species in Ukraine, Eur., Agr., Erg.
- Rubus laciniatus* Willd., Rosaceae, Shrub, Meso, Ken. (2013), unknown (Pyšek *et al.* 2012), Col., Erg.
- Rubus macropylloides* Weihe & Nees, Rosaceae, Shrub, Meso, Ken. (S.-E. of Ukraine, Ostapko, Boyko, Mosyakin 2010), C. Eur. (Krassovskaya 2001), Col., Kseno-Erg.
- Rudbeckia bicolor* Nutt., Asteraceae, Perenn., Meso, Ken., N. Am., Ergasphig., Erg.
- Rudbeckia hirta* L., Asteraceae, Ann. or bienn., Xero-meso, Ken., N. Am., Col., Erg.
- Rudbeckia laciniata* L., Asteraceae, Bienn., Hygro-meso, Ken., N. Am., Agr.-epoec., Erg.; invasive; transformer (Transcarpathia, Bukobinian Cis Carpathian).
- Rumex patientia* L., Polygonaceae, Perenn., Xero-meso, Ken., Med., Ergasphig., Erg.
- Sagittaria platyphylla* (Engelm.) J. G. Smith, Alismataceae, Perenn., Meso-hygro, Ken. (1982), N. Am., Ergasphig., Erg.
- Salix babylonica* L., Salicaceae, Tree, Meso, Ken., C.-E. As., Ephem., Erg.
- Salix fragilis* L., Salicaceae, Trees, Meso, Arch., As. min., Agr., Erg.; invasive; transformer (central regions of Ukraine).
- Salvia officinalis* L., Lamiaceae, Perenn., Xero-meso, Ken., Med., Ergasphig., Erg.
- Salvia splendens* Ker Gawl., Lamiaceae, Perenn., Xero, Ken., S. Am., Ephem., Erg.
- Sambucus racemosa* L., Sambucaceae, Shrub or tree, Meso, Ken., W. Eur., Ergasphig., Erg.
- Santolina chamaecyparissus* L., Asteraceae, Shrub, Meso-xero, Ken., Med., Ergasphig., Erg.
- Santolina virens* Mill., Asteraceae, Shrub, Meso-xero, Ken., S. Eur., Ergasphig., Erg.
- Saponaria officinalis* L., Caryophyllaceae, Perenn., Xero-meso, Ken., Med., Agr.-epoec., Erg.
- Satureja hortensis* L., Lamiaceae, Ann., Xero-meso, Ken., Med. (Protopopova 1991) or Cauc. As. Min. (Mayorov *et al.* 2012), Ephem., Erg.
- Scolymus maculatus* L., Asteraceae, Ann., Xero, Ken., Med., Ergasphig., Erg.
- Scorzonera hispanica* L., Asteraceae, Perenn., Xero-meso, Ken., Balcan., Ergasphig., Erg.
- Secale cereale* L., Poaceae, Ann., Xero-meso, Arch., E. Med., Ergasphig., Erg.
- Sedum spurium* M. Bieb., Crassulaceae, Perenn., Xero-meso, Ken., Cauc.-As. min., Ergasphig., Erg.

- Sempervivum tectorum* L., Crassulaceae, Perenn., Meso-xero, Ken., Middle Eur. (mount.), Ergasphig., Erg.
- Senecio cineraria* DC., Asteraceae, Perenn., Xero, Ken., W. Med., Agr., Erg.
- Setaria italica* (L.) P. Beauv., Poaceae, Ann., Xero-meso, Ken., As., Ergasphig., Erg.
- Sicyos angula* L., Cucurbitaceae, Ann., Meso, Ken., N. Am., Agr., Erg.
- Sida hermaphrodita* (L.) Rusby, Malvaceae, Perenn., Xero, Ken. (Crimea), N. Am., Ephem., Erg.
- Silene armeria* L., Caryophyllaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Silene pendula* L., Caryophyllaceae, Ann., Xero-meso, Ken., Med., Ergasphig., Erg.
- Silphium perfoliatum* L., Asteraceae, Perenn., Meso, Ken., N. Am., Col., Erg.
- Silybum marianum* (L.) Gaertn., Asteraceae, Ann. or bienn., Meso-xero, Ken., Med.-Iran.-Tur., Epoec., Erg.
- Sinapis alba* L., Brassicaceae, Ann., Xero-meso, Arch., Med.-Iran.-Tur., Epoec., Erg.
- Sisyrinchium septentrionale* Bicknell, Iridaceae, Perenn., Xero-meso, Ken., N. Am., Agr.-col., Erg.
- Solanum melongena* L., Solanaceae, Ann., Meso, Ken., As., Ergasphig., Erg.
- Solanum tuberosum* L., Solanaceae, Perenn., Meso, Ken., S. Am., Ergasphig., Erg.
- Solidago canadensis* L., Asteraceae, Perenn., Xero-meso, Ken., N. Am., Agr.-epoec., Erg.; invasive; transformer (forest and forest-steppe regions of Ukraine).
- Solidago graminifolia* (L.) Salisb., Asteraceae, Perenn., Meso, Ken., N. Am., Ergasphig., Erg.
- Solidago serotinoides* A. & D. Love, Asteraceae, Perenn., Meso, Ken., N. Am., Epoec., Erg.
- Sophora japonica* L., Fabaceae, Tree, Meso, Ken., E. As., Ergasphig., Erg.
- Sorghum ×alatum* Parodi, Poaceae, Perenn., Xero-meso, Ken., hybrid origin, Ergasphig., Erg.
- Sorghum saccharatum* (L.) Moench, Poaceae, Ann., Xero-meso, Ken., S. As., Ergasphig., Erg.
- Sorghum sudanense* (Piper) Stapf, Poaceae, Ann., Xero-meso, Ken., Afr., Ergasphig., Erg.
- Spartium junceum* L., Fabaceae, Shrub, Ken., Xero, Med., Col. Erg.
- Spinacia oleracea* L., Chenopodiaceae, Ann., Meso, Ken., E. Med., Ergasphig., Erg.
- Spiraea billardii* Hérincq, Rosaceae, Shrub, Xero-meso, Ken., hybrid origin, Ergasphig., Erg.
- Spiraea douglasii* Hook., Rosaceae, Shrub, Meso, Ken., N. Am., Ergasphig., Erg.
- Spiraea japonica* L.f., Rosaceae, Shrub, Meso, Ken., E. As., Ergasphig., Erg.
- Spiraea salicifolia* L., Rosaceae, Shrub, Xero-meso, Ken., Eur.-As., Ergasphig., Erg.
- Stachys byzantina* K. Koch, Lamiaceae, Perenn., Xero, Ken., Balcan.-As. Min., Ergasphig., Erg.
- Symporicarpos albus* (L.) S. F. Blake, Caprifoliaceae, Shrub, Meso-xero, Ken., N. Am., Ergasphig., Erg.
- Symphytum laevis* L., Asteraceae, Perenn., Meso, Ken., N. Am., Col., Erg.
- Symphytum lanceolatum* Willd., Asteraceae, Perenn., Meso, Ken., N. Am., Col., Erg.
- Symphytum novae-angliae* L., Asteraceae, Perenn., Meso, Ken., N. Am., Ergasphig., Erg.
- Symphytum novi-belgii* L., Asteraceae, Perenn., Meso, Ken., N. Am., Ergasphig., Erg.
- Symphytum ×salignus* Willd., Asteraceae, Perenn., Meso, Ken., hybrid origin, Ephem., Erg.; invasive.
- Symphytum ×versicolor* Willd., Asteraceae, Perenn., Meso, Ken., hybrid origin, Ephem., Erg.
- Sympyrum asperum* Lepech., Boraginaceae, Perenn., Meso, Ken., Cauc.-E. Med., Ergasphig., Erg.
- Sympyrum caucasicum* M. Bieb., Boraginaceae, Perenn., Meso, Ken., Cauc., Col., Erg.
- Sympyrum peregrinum* Ledeb., Boraginaceae, Perenn., Meso, Ken., Cauc., Ergasphig., Erg.
- Syringa vulgaris* L., Oleaceae, Shrub, Xero-meso, Ken., S. Eur., Col., Erg.
- Tagetes erecta* L., Asteraceae, Ann., Meso, Ken., N. Am., Ergasphig., Erg.
- Tamarix ramosissima* Ledeb., Tamaricaceae, Shrub or tree, Meso-xero, Ken., Middle As., Agr., Erg.
- Tamarix smyrnensis* Bunge, Tamaricaceae, Shrub, Meso-xero, Ken., As. Min. and Middle, Ergasphig., Erg.
- Tamarix tetrandra* Pall. ex M. Bieb., Tamaricaceae, Shrub, Meso-xero, Ken. (Cis Black Sea Region), E. Med.-As. Min., Ergasphig., Erg.
- Thladiantha dubia* Bunge, Cucurbitaceae, Perenn., Meso, Ken., S.-E. As., Col., Erg.
- Thlaspi perfoliatum* L., Brassicaceae, Ann., Meso, Ken., Med., Ergasphig., Erg.
- Thuja occidentalis* L., Cupressaceae, Tree, Meso, Ken. N. Am., Ergasphig., Erg.
- Toxicodendron radicans* (L.) O. Kunze, Anacardiaceae, Liane, Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Tradescantia virginiana* L., Commelinaceae, Perenn., Hygro-meso, Ken., N. Am., Ergasphig., Erg.
- Tragopogon porrifolius* L., Asteraceae, Perenn., Xero-meso, Ken., Med., Ergasphig., Erg.
- Trifolium hybridum* L., Fabaceae, Ann. or bienn., Meso, Ken., unknown, Ergasphig., Erg.
- Trifolium incarnatum* L., Fabaceae, Ann., Meso, Ken., W. Med., Ergasphig., Erg.
- Trifolium resupinatum* L., Fabaceae, Ann., Xero-meso, Ken., Med., Ephem., Erg.-kseno.
- Trifolium sativum* (Schreb.) Crome, Fabaceae, Perenn., Xero-meso, Ken., W.-C. Eur., Ergasphig., Erg.
- Trigonella caerulea* (L.) Ser., Fabaceae, Ann., Xero-meso, Ken., Med., Ephem., Erg.-kseno.
- Trigonella foenum-graecum* L., Fabaceae, Ann., Xero-meso, Ken., W. As., Ergasphig., Erg.
- Triticum aestivum* L., Poaceae, Ann., Xero-meso, Ken., As., Ergasphig., Erg.
- Triticum durum* Desf., Poaceae, Ann., Xero-meso, Ken., As., Ergasphig., Erg.
- Tropaeolum majus* L., Tropaeolaceae, Ann., Meso, Ken., S. Am., Ergasphig., Erg.
- Tropaeolum minus* L., Tropaeolaceae, Ann., Meso, Ken., S. Am., Ergasphig., Erg.
- Tulipa sylvestris* L., Liliaceae, Perenn., Xero-meso, Ken. (2013), Med., Col., Erg.
- Ulex europaeus* L., Fabaceae, Shrub, Xero, Ken., C.-W. Eur., Ergasphig., Erg.
- Ulmus pumila* L., Ulmaceae, Tree, Meso-xero, Ken., As., Epoec., Erg.
- Verbesina encelioides* (Cav.) Benth. & Hook. f. ex A. Gray, Asteraceae, Ann., Xero, Ken. (1934), N. Am., Epoec., Erg.
- Veronica filiformis* Smith, Scrophulariaceae, Ann.-bienn., Xero-meso, Ken., Med. (Pyšek *et al.* 2012), Ephem., Erg.-kseno.
- Viburnum tinus* L., Viburnaceae, Shrub, Xero-meso, Ken., Med., Agr. (Crimea), Erg.
- Vicia faba* L., Fabaceae, Ann., Meso, Ken., unknown, Ergasphig., Erg.
- Vicia sativa* L., Fabaceae, Ann., Meso, Ken., Med.-Afr.-As (Pyšek *et al.* 2012), Ephem., Erg.

- Vinca major* L., Apocynaceae, Perenn., Xero-meso, Ken. (Crimea), Med., Ergasphig., Erg.
- Viola nemausensis* Jord., Violaceae, Ann., Xero-meso, Ken., unknown, Ergasphig., Erg.
- Viola ×wittrockiana* Gams ex Hegi, Violaceae, Perenn., Meso, Ken., hybrid origin, Ergasphig., Erg.
- Vitex agnus-castus* L., Verbenaceae, Shrub, Xero-meso, Ken., Eur.-As., Ergasphig., Erg.
- Vitis amurensis* Rupr., Vitaceae, Liana, Meso, Ken., Far E., Col., Erg.
- Vitis labrusca* L., Vitaceae, Liana, Xero-meso, Ken., N. Am., Ergasphig., Erg.
- Vitis vinifera* L., Vitaceae, Liana, Meso-xero, Ken., unknown (Pyšek *et al.* 2012), Ergasphig., Erg.
- Vitis vulpina* L., Vitaceae, Liana, Xero-meso, Ken., N. Am., Col., Erg.
- Zea mays* L., Poaceae, Ann., Xero-meso, Ken., C.-S. Am., Ergasphig., Erg.
- Zelkova carpinifolia* (Pall.) K. Koch, Ulmaceae, Tree, Meso, Ken. (Crimea 1987), Cauc., Col., Erg.
- Zinnia elegans* Jacq., Asteraceae, Ann., Meso, Ken., N. Am., Ergasphig., Erg.
- Zizania aquatica* L., Poaceae, Ann., Hydro, Ken., N. Am., Col., Erg.
- Zizania latifolia* (Griseb.) Stapf, Poaceae, Perenn., Hydro, Ken., E. As., Agr., Erg.

Explanations: **life forms**, Ann. – annual, Bien. – biennial, Ann.-bienn. – annual or biennial, Perenn. – perennial; **hydromorphe**, Hydro – hydrophytes, Hygro – hygrophytes, Hygro-meso – hygro-mesophytes, Meso – mesophytes, Meso-hydro – meso-hygrophyte, Meso-xero – meso-xerophytes, Xero – xerophytes, Xero-meso – xero-mesophytes; **chronological elements**, Arch. – archaeophytes, Ken. – kenophytes; **degree of naturalization**, Agr. – agriophytes, Agr.-epoec. – agrio-epoecophytes, Col. – colonophytes, Ephem. – ephemeroophytes, Epoec. – epoecophytes, Ergasphig. – ergasiophyophytes; **origin**, Afr. – African, Am. – American, As. – Asia, As. Min. – Asia Minor, As. Min.-Middle As. – Asia Minor-Middle Asia, Balk. – Balcan, Balk.-C. As. – Balcan-Central Asian, C. Am. – Central American, C. As. – Central Asian, C. Eur. – Central European, Cauc. – Caucasic, Cauc.-As. Min. – Caucasic-Asia Minor, Cauc.-As. Min.-Med. – Caucasic-Asia Minor –Mediterraneum, Cauc.-E. Med. – Caucasic-Eastern Mediterraneum, Cauc.-Eur. – Caucasic-European, E. As. – Eastern Asia, Eur. – European, Eur.-As. – European-Asian, Eur.-Cauc. – European-Caucasic, E. Med. – Eastern Mediterraneum, E. Med.-As. Min. – Eastern Mediterraneum-Asia Minor, Far E. – Far East, Iran-Tur. – Iran-Turanian, Med. – Mediterraneum, Med.-Afr.-As. – Mediterraneum-African-Asian, Med.-As. – Mediterraneum-Asian, Med.-Iran.-Tur. – Mediterraneum-Iran-Turanian, Med.-Middle As. – Mediterraneum-Middle Asian, Middle As. – Middle Asia, Middle As. – Middle Asian, Middle Eur. – Middle European, Middle Eur.-Balk. – Middle European-Balkan, N. Afr. – Northern African, N.-W. Afr. – Northern-Western African, N. Am. – North American, N. Pann. – Northern Pannonian, S. Cis Black Sea – South Cis Black Sea, S. Am. – South American, S. As. – South Asia, S.-E. As. – South-Eastern Asia, S.-E. Transcauc. – South-Eastern Trasncaucasian, S.-Eur. – South European, Sib. – Siberian, sub.-Alp. – subalpic, Trop. Am. – Tropical American, Tur. – Turanian, W. As. – West Asia, W. As. Min. – Western Asia Minor, W.-C. Eur. – West-Central European, W. Med. – West Mediterraneum, W. Eur. – West European; **way of distribution**, Erg. – ergasiophytes, Erg.-kseno. – ergasio-ksenophytes